

# Weakly coupled two slow- two fast systems, folded node and mixed mode oscillations

**M. Krupa<sup>1</sup>, B. Ambrosio, M.A. Aziz-Alaoui<sup>2</sup>**

<sup>1</sup> INRIA Paris-Rocquencourt Research Center, Domaine de Voluceau BP 105, 78153 Le Chesnay cedex, France

<sup>2</sup> Normandie Univ, France; ULH, LMAH, F-76600 Le Havre; FR CNRS 3335, 25 rue Philippe Lebon 76600 Le Havre, France.

E-mail: benjamin.ambrosio@univ-lehavre.fr

**Abstract.** We study Mixed Mode Oscillations (MMOs) in systems of two weakly coupled slow/fast oscillators. We focus on the existence and properties of a folded singularity called FSN II that allows the emergence of MMOs in the presence of a suitable global return mechanism. As FSN II corresponds to a transcritical bifurcation for a desingularized reduced system, we prove that, under certain non-degeneracy conditions, such a transcritical bifurcation exists. We then apply this result to the case of two coupled systems of FitzHugh-Nagumo type. This leads to a non trivial condition on the coupling that enables the existence of MMOs.

AMS classification scheme numbers: 34C15, 34D15, 34K18.

## 1. Introduction

*Canard phenomenon*, known also as *canard explosion* is a transition from a small, Hopf type oscillation to a relaxation oscillation, occurring upon variation of a parameter. This transition has been first found in the context of the van der Pol equation [2], and soon after in numerous models of phenomena occurring in engineering and in chemical reactions [4]. A common feature of all these models is the presence of time scale separation (one slow, one fast variable). A particular feature of canard explosion is that it takes place in a very small parameter interval. For the van der Pol system, where the ratio of the timescales is given by a small parameter  $\varepsilon$ , the width of this parameter interval can be estimated by  $O(\exp(-c/\varepsilon))$ , where  $c > 0$  is a fixed positive constant. The transition occurs through a sequence of *canard cycles* whose striking feature is that they contain segments following closely a stable slow manifold and subsequently an unstable slow manifold.

The work on canard explosion led to investigations of slow/fast systems in three dimensions, with two slow and one fast variables. In the context of these systems the concept of a *canard solution* or simply *canard* has been introduced, as a solution passing from a stable to an unstable slow manifold [1, 12, 16]. Canards arise near so called folded singularities of which the best known is folded node [1, 16] and [18]. Unlike in systems with one slow variable, canards occur robustly (in  $O(1)$  parameter regions) in systems with two slow variables. Related to canards are Mixed Mode Oscillations (MMOs), which are trajectories that combine small oscillations and large oscillations of relaxation type, both recurring in an alternating manner. Recently there has been a lot of interest in MMOs that arise due to the presence of canards, starting with the work of Milik, Szmolyan, Loeffelmann and Groeller [12]. The small oscillations arise during the passage of the trajectories near a folded node or a related folded singularity. The dynamics near the folded singularity is transient, yet recurrent: the trajectories return to the neighborhood of the folded singularity by way of a global return mechanism [3].

The setting of folded node combined with a global return mechanism, elucidated in [3], led to the explanations of MMO dynamics found in applications [15, 14, 8, 17]. A shortcoming of the folded node setting is the lack of connection to a Hopf bifurcation, which seems to play a prominent role in many MMOs. This led to the interest in another, more degenerate, folded singularity, known as *Folded Saddle Node of type II* (FSNII), originally introduced in [12] and recently analyzed in some detail by Krupa and Wechselberger [11]. Guckenheimer [10] studied a very similar problem in the parameter regime yet closer to the Hopf bifurcation, calling it *singular Hopf bifurcation*. The transition between the two settings was studied in [6]. Another interesting singularity, that was mentioned in [11] and can lead to rich families of MMOs, without the presence of a Hopf bifurcation, is *Folded Saddle Node of type I* (FSN I). For a more comprehensive overview we refer the reader to the recent review article [7]. The theory of two slow and one fast variables has recently been generalized by Wechselberger [20] to arbitrary finite dimensions.

In this article we study systems of two weakly coupled slow/fast oscillators. We assume that in the absence of the coupling one of the oscillators is undergoing a canard explosion

and the other is in general position. We show that turning on the weak coupling leads to the occurrence of MMOs. We focus on the very interesting case of FSN II, which, in the uncoupled case, corresponds to one of the oscillators undergoing a canard explosion, while the other is at a stable equilibrium. As elaborated in [3] canard induced MMOs arise through a combined presence of a folded singularity and suitable return mechanism. In this paper we focus on the existence and properties of a folded singularity leaving the study of the return mechanism for future investigations.

The article is organized as follows. We start with a background, Section 2, in which we explain the very standard case of MMOs in the context of two slow and one fast variables and subsequently the case of two slow and two fast variables with a simple fold line. This material is included for completeness and presented in such a way that the coupled oscillator case becomes a simple corollary. In Section 3, we treat the general case of coupled oscillators and in Section 4, we consider the example of two coupled FitzHugh-Nagumo systems.

## 2. Background

### 2.1. The basic case of one fast and two slow variables

We consider a system of the form

$$\varepsilon \dot{x} = f(x, y), \quad (1)$$

$$\dot{y} = g(x, y), \quad x \in \mathbb{R}, y = (y_1, y_2) \in \mathbb{R}^2, g = (g_1, g_2). \quad (2)$$

The functions  $f$  and  $g$  may depend on  $\varepsilon$  and other parameters, but we suppress this dependence for simplicity of notation. The associated reduced system is

$$0 = f(x, y), \quad (3)$$

$$\dot{y} = g(x, y), \quad x \in \mathbb{R}, y = (y_1, y_2) \in \mathbb{R}^2. \quad (4)$$

Let  $S_0$  denote the curve defined by  $0 = f(x, y)$ . Non-hyperbolic points correspond to the points on  $S_0$  for which  $f_x = 0$  (we use the notation  $f_\xi = \partial f / \partial \xi$ ). At such points the equation  $0 = f(x, y)$  cannot be solved for  $x$  as a function of  $y$ . Suppose  $(0, 0)$  is a non-hyperbolic point. We make a non-degeneracy assumption  $f_{xx}(0) \neq 0$ . In order to obtain an explicit equation for the slow flow, we try to solve  $0 = f(x, y)$  for  $y_1$  or  $y_2$ . It is convenient to first change the variables to simplify the process of finding such a solution. We begin with a change of variables of the form  $x \rightarrow x + \eta(y)$ ,  $\eta(0, 0) = 0$ . In the new variables, with additional scaling,  $f$  has the form

$$f(x, y_1, y_2) = f(0, y) + x^2 + O(x^2|y|, x^3).$$

We make a non-degeneracy assumption  $f_y(0, 0) \neq (0, 0)$ . Without loss of generality (WLOG), we can assume that  $f_{y_1}(0, 0) \neq 0$ . We can now introduce a new coordinate  $\tilde{y}_1 = -f(0, y)$ , and immediately drop the tilde, to simplify the notation. In the new coordinates  $f$  has the form

$$f(x, y_1, y_2) = -y_1 + x^2 + O(x^2|y|, x^3).$$

The transformations we made may be just valid locally, i.e. only in a small neighborhood of the non-hyperbolic point. Now  $S_0$  is (locally) represented as a graph:

$$y_1 = x^2(1 + O(y_2, x)). \quad (5)$$

Differentiating (5) we get

$$\dot{y}_1 = 2x(1 + O(y_2, x))\dot{x} + O(x^2). \quad (6)$$

Substituting (6) into (4) we get

$$2x(1 + O(y_2, x))\dot{x} = g_1(x, x^2(1 + O(y_2, x)), y_2) + O(x^2) \quad (7)$$

$$\dot{y}_2 = g_2(x, x^2(1 + O(y_2, x)), y_2). \quad (8)$$

Now it is clear that (7) has a singular at the fold as long as  $g_1(0, 0, y_2) \neq 0$ . Points on the fold line for which  $g_1(0, 0, y_2) = 0$  are called folded singularities. We would like to understand the flow near such points better and to this end we apply a singular time rescaling to (7)-(8) by multiplying the right-hand side (RHS) by the factor  $2x(1 + O(y_2, x))$  and canceling it in (7). This leads to the following system:

$$\dot{x} = g_1(x, x^2(1 + O(y_2, x)), y_2) + O(x^2) \quad (9)$$

$$\dot{y}_2 = 2x(1 + O(y_2, x))g_2(x, x^2(1 + O(y_2, x)), y_2). \quad (10)$$

We refer to (9)-(10) as the desingularized system. Note that folded singularities correspond to equilibria of (9)-(10) with  $x = 0$ . Further note that the trajectories of (7)-(8) and (9)-(10) restricted to the half plane  $x > 0$  differ only by time parametrization but are the same as sets. The trajectories in the half-plane  $x < 0$  are the same as sets but have the opposite direction of time. The flow near folded singularities is determined by the linearization of (9)-(10) at folded singularities, which is given by the Jacobian matrix:

$$\begin{pmatrix} g_{1,x}(0, 0, y_2) & g_{1,y_2}(0, 0, y_2) \\ 2g_2(0, 0, y_2) & 0 \end{pmatrix}. \quad (11)$$

Folded singularities are classified according to the type of the corresponding equilibrium of (9)-(10). Canards arise near folded saddles and folded nodes and small oscillations are associated with folded nodes, see the work of Benoît [1], and Szmolyan and Wechselberger [16]. Folded node, which is the only generic folded singularity whose dynamics is accompanied by small oscillations, occurs when the corresponding equilibrium of (9)-(10) has two real positive eigenvalues.

## 2.2. Folded singularities

We now consider the case when  $(f, g)$  (the RHS of (1)-(2)) depend on a regular parameter and  $g_2(0, 0, y_2)$  passes through 0 when this parameter is varied. This passage corresponds to change of sign of the determinant of the Jacobian (11), from negative to positive or vice-versa. This, in turn, means that the eigenvalues change from one positive and one negative (negative determinant) to two positive (positive determinant), i.e. from saddle to node. It follows that this transition corresponds to the onset of small oscillations. Hence the onset of

small oscillations is a consequence of the passage of  $g_2(0, 0, y_2)$  through 0. This transition is called Folded Saddle-Node of type II (FSN II).

As finding FSN II in coupled oscillator systems is the focus of this article we review some of the features of this bifurcation, referring the reader to [11] for details. An important feature of FSN II is that it corresponds to the passage of a true equilibrium of (3)-(4) through the fold line. We assume WLOG that FSN II corresponds to the point  $(x, y, \lambda) = (0, 0, 0)$ , where  $\lambda$  is the regular parameter. Suppose that FSN II is *non-degenerate*, that is  $(x, y, \lambda) = (0, 0, 0)$  corresponds to a transcritical bifurcation of (9)-(10). Then, for  $\lambda = 0$ , there is a folded singularity satisfying  $g_2(0, 0, 0) = 0$  and, for  $\lambda \neq 0$  but close to 0 there is a point close to the origin, such that  $g_1(x, x^2 + O(x^3), y_2) = g_2(x, x^2 + O(x^3), y_2) = 0$ , i.e. a true equilibrium of (3)-(4) near the fold line. For the original system (1)-(2) one can prove that this corresponds to a so called singular Hopf bifurcation [10], and, in a different regime of the parameter  $\lambda$ , to a delayed Hopf bifurcation [11]. These two bifurcations lead to the existence of small oscillations and thus enable the existence of MMOs. We note here that the interesting case of FSN II is when the equilibrium is stable on the stable slow manifold and a saddle on the unstable slow manifold.

For folded nodes, not close to either FSN II or to Folded Saddle Node of type I (FSN I), defined by  $g_{1,y_2} = 0$ , small oscillations are extremely small, which means that they cannot be seen, except with very detailed numerics, see [3]. Hence the most interesting cases occur near FSN I or FSN II.

### 2.3. Canonical system in two fast and two slow dimensions

We consider a system with two fast and two slow variables

$$\varepsilon \dot{x} = f(x, y, \varepsilon), \tag{12}$$

$$\dot{y} = g(x, y, \varepsilon), \quad x = (x_1, x_2) \in \mathbb{R}^2, \quad y = (y_1, y_2) \in \mathbb{R}^2, \tag{13}$$

$f = (f_1, f_2)$ ,  $g = (g_1, g_2)$ . Recall that in the fast formulation (12)-(13) has the form

$$x' = f(x, y, \varepsilon), \tag{14}$$

$$y' = \varepsilon g(x, y, \varepsilon). \tag{15}$$

Recall also that the reduced problem has the form

$$0 = f(x, y, 0), \tag{16}$$

$$\dot{y} = g(x, y, 0). \tag{17}$$

and the layer problem has the form

$$x' = f(x, y, \varepsilon), \tag{18}$$

$$y' = 0. \tag{19}$$

We consider a point on the fold curve (WLOG we assume that this point is the origin  $(0, 0)$ ).

Let  $\Phi : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  be the map defined by

$$\Phi(x, y) = \begin{pmatrix} f_1(x, y) \\ f_2(x, y) \\ \det(f_x)(x, y) \end{pmatrix}. \tag{20}$$

The fold curve is defined by the condition  $\Phi = 0$ . We consider a point on the fold line (WLOG we assume that this point is the origin  $(0, 0)$ ). We now state a few conditions which assure that the fold curve is *simple*; naturally the first condition is that the linearization of layer system has a simple eigenvalue 0. More specifically, our first assumption is as follows:

$f_x(0, 0)$  has one simple 0 eigenvalue and one simple eigenvalue in the left half-plane. (21)

Recall that in Section 2.1 we made additional assumptions, namely  $f_{xx}(0, 0) \neq 0$  and  $f_y(0, 0) \neq (0, 0)$ . Here we generalize these conditions in the following way. Let  $(v_1, v_2)$  denote the nullvector (the eigenvector of 0) and  $(w_1, w_2)$  the eigenvector of the negative eigenvalue. Our additional non-degeneracy conditions are as follows:

$$\text{either } f_{y_1}(0, 0) \text{ or } f_{y_2}(0, 0) \text{ is independent of } \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} \quad (22)$$

$$\kappa = (\det(f_x))_x(0, 0) \cdot (v_1, v_2) \neq 0. \quad (23)$$

We now have the following proposition.

**Proposition 1** *We assume that  $(0, 0)$  is on the fold curve, i. e.  $\Phi(0, 0) = 0$ , and that conditions (21)-(23) are satisfied. WLOG we assume that  $f_{y_1}(0, 0)$  is independent of  $(w_1, w_2)$ . Then there exists a (locally) unique solution of the equation  $\Phi = 0$ . Moreover, this solution can be expressed as  $(\psi_1(y_2), \psi_2(y_2), \psi_3(y_2), y_2)$ , where  $\psi = (\psi_1, \psi_2, \psi_3) : \mathbb{R} \rightarrow \mathbb{R}^3$  is a smooth function, with  $\psi(0) = (0, 0, 0)$ .*

**Proof** To apply the implicit function theorem we verify that the  $3 \times 3$  matrix  $M = \Phi_{(x_1, x_2, y_1)}(0, 0)$  is non singular. We do this by finding three independent vectors in the image of  $M$ . First note that  $M(v_1, v_2, 0) = (0, 0, \kappa)$  and that the first two coordinates of  $M(w_1, w_2, 0)$  are a multiple of  $(w_1, w_2)$ . Note also that the first two coordinates of  $M(0, 0, 1)$  are equal to  $f_{y_1}(0, 0)$ . By assumption  $f_{y_1}(0, 0)$  is independent of  $(w_1, w_2)$ , hence the vectors  $M(v_1, v_2, 0)$ ,  $M(w_1, w_2, 0)$  and  $M(0, 0, 1)$  are linearly independent. It follows that the equation  $\Phi = 0$  can be solved by the implicit function theorem, giving  $(x_1, x_2, y_1)$  as a function of  $y_2$ .

In the remainder of this section we will assume that the fold curve has been transformed to the coordinate line  $(0, 0, 0, y_2)$  and that the fast system has been diagonalized. More specifically, we assume that  $f$  satisfies the following conditions

$$f(0, 0, 0, y_2, 0) = 0, \quad \frac{\partial f_1}{\partial x_1}(0, 0, 0, y_2, 0) = 0, \quad \frac{\partial f_2}{\partial x_2}(0, 0, 0, y_2, 0) = \lambda(y_2) < 0, \quad (24)$$

$$\frac{\partial f_1}{\partial x_2}(0, 0, 0, y_2, 0) = \frac{\partial f_2}{\partial x_1}(0, 0, 0, y_2, 0) = 0. \quad (25)$$

Note that, since  $(v_1, v_2) = (1, 0)$  and  $(w_1, w_2) = (0, 1)$  the non degeneracy conditions (22)-(23) reduce to

$$\frac{\partial f_1}{\partial y_1}(0, 0, 0, y_2, 0) \neq 0, \quad \frac{\partial^2 f_1}{\partial x_1^2}(0, 0, 0, y_2, 0) \neq 0. \quad (26)$$

WLOG we assume

$$\frac{\partial f_1}{\partial y_1}(0, 0, 0, y_2, 0) = -1, \quad \frac{\partial^2 f_1}{\partial x_1^2}(0, 0, 0, y_2, 0) = 1.$$

We can now expand  $f$  in Taylor series:

$$f(x, y, \varepsilon) = \begin{pmatrix} -y_1 + x_1^2 + O(x_1 x_2, x_2^2) + O(3) + O(\varepsilon) \\ \lambda(y)x_2 + O(y_1, \|x\|^2) + O(\varepsilon) \end{pmatrix}. \quad (27)$$

The defining conditions of the slow manifold are  $f(x, y, 0) = 0$ , or

$$0 = -y_1 + x_1^2 + O(x_1 x_2, x_2^2) + O(3), \quad (28)$$

$$0 = \lambda(y)x_2 + O(y_1, \|x\|^2) \quad (29)$$

From (29) we get, by the implicit function theorem,  $x_2 = p(x_1, y)$ , with  $p(x_1, y) = O(y_1, x_1^2)$ . Plugging into (28) we get the usual condition  $y_1 = x_1^2 + O(x_1^3)$ . Following the approach of Section 2.1 we substitute  $y_1 = x_1^2 + O(x_1^3)$  into (17) getting

$$(2x_1 + O(x_1^2))\dot{x}_1 = g_1(x_1, p(x_1, x_1^2 + O(x_1^3)), y_2, x_1^2 + O(x_1^3), y_2, 0), \quad (30)$$

$$\dot{y}_2 = g_2(x_1, p(x_1, x_1^2 + O(x_1^3)), y_2, x_1^2 + O(x_1^3), y_2, 0). \quad (31)$$

Finally we get the desingularized equation as follows:

$$\dot{x}_1 = g_1(x_1, p(x_1, x_1^2 + O(x_1^3)), y_2, x_1^2 + O(x_1^3), y_2, 0), \quad (32)$$

$$\dot{y}_2 = g_2(x_1, p(x_1, x_1^2 + O(x_1^3)), y_2, x_1^2 + O(x_1^3), y_2, 0)(2x_1 + O(x_1^2)) \quad (33)$$

Folded singularities are points  $(0, y_2)$  with  $y_2$  satisfying  $g_1(0, 0, 0, y_2, 0) = 0$ , or, equivalently, equilibrium points of (32)-(33) on the fold line, i.e. satisfying  $x_1 = 0$ . The type of folded singularity is determined by the Jacobian of

$$\begin{pmatrix} g_{1,x_1}(0, 0, 0, y_2) & g_{1,y_2}(0, 0, 0, y_2) \\ 2g_2(0, 0, 0, y_2) & 0 \end{pmatrix}. \quad (34)$$

Folded saddle nodes (FSN I and FSN II) can arise in the context of systems with two slow and two fast dimensions in an analogous way as described in Section 2.2 for systems with two slow and one fast dimensions, and, in the same manner, correspond to the onset of MMOs.

### 3. Coupled oscillator system

#### 3.1. Introduction of the system

We consider a system of coupled oscillators in the following form

$$\varepsilon \dot{x}_1 = -y_1 + F_1(x_1) + \nu H_1^f(x, y) \quad (35)$$

$$\varepsilon \dot{x}_2 = -y_2 + F_2(x_2) + \nu H_2^f(x, y) \quad (36)$$

$$\dot{y}_1 = G_1(x_1, y_1, c_1) + \nu H_1^s(x, y) \quad (37)$$

$$\dot{y}_2 = G_2(x_2, y_2, c_2) + \nu H_2^s(x, y) \quad x = (x_1, x_2) \in \mathbb{R}^2, y = (y_1, y_2) \in \mathbb{R}^2 \quad (38)$$

The parameters  $\varepsilon$  and  $\nu$  are the singular parameter and the coupling parameter, respectively, and are considered to be small. The parameters  $c_1, c_2$  controls the state of the uncoupled oscillators (moves the nullclines). The coupling functions  $H^f : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ ,  $H^s : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  and  $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  are defined by  $H^f = (H_1^f, H_2^f)$ , by  $H^s = (H_1^s, H_2^s)$  and  $F = (F_1, F_2)$ ,

respectively. We assume that  $y_j = F_j(x)$  are  $S$  shaped curves. Written on the fast time scale (35)-(38) has the form:

$$x'_1 = -y_1 + F_1(x_1) + \nu H_1^f(x, y) \quad (39)$$

$$x'_2 = -y_2 + F_2(x_2) + \nu H_2^f(x, y) \quad (40)$$

$$y'_1 = \varepsilon G_1(x_1, y_1, c_1) + \nu H_1^s(x, y) \quad (41)$$

$$y'_2 = \varepsilon G_2(x_2, y_2, c_2) + \nu H_2^s(x, y) \quad (42)$$

We now find the conditions for the existence of a simple fold curve, as in Section 2.3. Let  $\Phi$  defined as in section 2.3. First note that the critical manifold of (35)-(38) (the constraint manifold of (35)-(38) with  $\varepsilon = 0$ ), is defined by

$$-y_j + F_j(x) + \nu H_j^f(x, y) = 0, \quad j = 1, 2. \quad (43)$$

The linearization of the RHS of (35)-(36) is given by

$$\begin{pmatrix} F'_1(x_1) + \nu H_{1,x_1}^f & \nu H_{1,x_2}^f \\ \nu H_{2,x_1}^f & F'_2(x_2) + \nu H_{2,x_2}^f \end{pmatrix} \quad (44)$$

where we assume that  $(x, y)$  is on the critical manifold, i.e. satisfies (43). The determinant of the matrix in (44) is as follows:

$$F'_1(x_1)F'_2(x_2) + \nu(F'_1(x_1)H_{2,x_2}^f + F'_2(x_2)H_{1,x_1}^f) + \nu^2 \det H_x^f. \quad (45)$$

**Proposition 2** *Let  $(x_0, y_0)$  satisfy  $\Phi(x_0, y_0) = 0$  for  $\nu = 0$ . We assume that*

$$F'_1(x_{0,1}) = 0 \text{ and } F'_2(x_{0,2}) < 0 \text{ as already specified in (24).}$$

*And that,*

$$F''_1(x_{0,1}) \neq 0.$$

*Then, there exists a (locally) unique solution of the equation  $\Phi = 0$ . This solution can be expressed as  $(\psi_1(y_2), \psi_2(y_2), \psi_3(y_2), y_2)$ , where  $\psi = (\psi_1, \psi_2, \psi_3) : \mathbb{R} \rightarrow \mathbb{R}^3$  is a smooth function, with  $\psi(y_{2,0}) = (x_{1,0}, x_{2,0}, y_{1,0})$ . Furthermore, the parametrization of the fold curve depends also smoothly on the parameters  $\nu$ .*

### Proof

We apply Proposition 1. For this we check the conditions (21), and (22)-(23). When  $F'_1(x_{0,1}) = 0$  and  $F'_2(x_{0,2}) \neq 0$ , (21) hold (note that one could choose  $F'_1(x_{0,1}) \neq 0$  and  $F'_2(x_{0,2}) = 0$ ). We now have, for  $\nu = 0$ ,  $(v_1, v_2) = (1, 0)$  and  $(w_1, w_2) = (0, 1)$ . It is now easy to verify that (22) holds automatically and (23) is equivalent to  $F''_1(x_{0,1}) \neq 0$ . By continuity, the result remains valid in a neighborhood of  $\nu = 0$ . Note that, because the fast system does not depend on parameters  $c_1, c_2$ , the fold curve also does not.

### 3.2. Folded singularities and their nature

Let  $x_1^* = \psi_1(y_2, \nu)$ ,  $x_2^* = \psi_2(y_2, \nu)$  and  $y_1^* = \psi_3(y_2, \nu)$ . We transform the fold curve to the line  $(0, 0, 0, y_2)$ , using the following change of variables:  $\tilde{x}_1 = x_1 - x_1^*$ ,  $\tilde{x}_2 = x_2 - x_2^*$ ,  $\tilde{y}_1 = y_1 - y_1^*$ . We omit the tilde to simplify the notation. System (35)-(38) becomes

$$\varepsilon \dot{x}_1 = -y_1 + f_1(x_1, y_2) + \nu h_1^f(x, y) + O(\varepsilon) \quad (46)$$

$$\varepsilon \dot{x}_2 = f_2(x_2, y_2) + \nu h_2^f(x, y) + O(\varepsilon) \quad (47)$$

$$\begin{aligned} \dot{y}_1 &= G_1(x_1 + x_1^*, y_1 + y_1^*, c_1) + \nu H_1^s(x + x^*, y_1 + y_1^*, y_2) \\ &\quad - \frac{dy_1^*}{dy_2} (G_2(x_2 + x_2^*, y_2, c_2) + \nu H_2^s(x + x^*, y_1 + y_1^*, y_2)) \end{aligned} \quad (48)$$

$$\dot{y}_2 = G_2(x_2 + x_2^*, y_2, c_2) + \nu H_2^s(x + x^*, y_1 + y_1^*, y_2), \quad (49)$$

where,

$$f_1(x_1, y_2) = F_1'(x_1^*)x_1 + F_1''(x_1^*)\frac{1}{2}x_1^2 + O(x_1^3),$$

$$f_2(x_2, y_2) = F_2'(x_2^*)x_2 + F_2''(x_2^*)\frac{1}{2}x_2^2 + O(x_2^3),$$

and,

$$\begin{aligned} h_1^f(x, y) &= H_{1,x_1}^f(x^*, y_1^*, y_2)x_1 + H_{1,x_2}^f(x^*, y_1^*, y_2)x_2 + H_{1,y_1}^f(x^*, y_1^*, y_2)y_1 + O(\|(x, y_1)\|^2) \\ h_2^f(x_2, y_2) &= H_{2,x_1}^f(x^*, y_1^*, y_2)x_1 + H_{2,x_2}^f(x^*, y_1^*, y_2)x_2 + H_{2,y_1}^f(x^*, y_1^*, y_2)y_1 + O(\|(x, y_1)\|^2). \end{aligned}$$

Our goal is to arrive at the desingularized system (32)-(33). The first step is to diagonalize the linear part of the fast subsystem (46)-(47). More precisely, we change coordinates so that the RHS of (46)-(49) is transformed to the form (27). Let

$$\begin{pmatrix} 1 \\ v(y_2) \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} w(y_2) \\ 1 \end{pmatrix} \quad (50)$$

be the eigenvectors of the Jacobian (44) at the points  $(0, 0, 0, y_2)$  on the fold line, corresponding to the eigenvalues 0 and  $\lambda(y_2) < 0$ , respectively. The diagonalizing transformation has the form

$$\tilde{x} = P(y)^{-1}x,$$

where the columns of the matrix  $P$  are the eigenvectors (50). In the new variables system (46)-(49) becomes (tilde is omitted)

$$\varepsilon \dot{x}_1 = \frac{1}{1 - vw} \left( -y_1 + \frac{K}{2}x_1^2 + \nu(O(y_1)) + O(x_1x_2, x_2^2) + O(\varepsilon) \right) \quad (51)$$

$$\varepsilon \dot{x}_2 = \lambda(y_2)x_2 + \nu(O(y_1)) + O(\|x\|^2) + O(\varepsilon) \quad (52)$$

$$\begin{aligned} \dot{y}_1 &= G_1(x_1 + x_2w(y_2) + x_1^*, y_1 + y_1^*, c_1) + \nu H_1^s(P(y_2)x + x^*, y_1 + y_1^*, y_2) \\ &\quad - \frac{dy_1^*}{dy_2} \left( G_2(x_2 + x_1v(y_2) + x_2^*, y_2, c_2) + \nu H_2^s(P(y_2)x + x^*, y_1 + y_1^*, y_2) \right) \end{aligned} \quad (53)$$

$$\dot{y}_2 = G_2(x_2 + x_1v(y_2) + x_2^*, y_2, c_2) + \nu H_2^s(P(y_2)x + x^*, y_1 + y_1^*, y_2), \quad (54)$$

where,

$$K = F_1''(x_1^*) + O(\nu). \quad (55)$$

Note that the fast subsystem (51)-(52) is now as specified in (27). Applying the procedure described in Section 2.3 we obtain the reduced system

$$(Kx_1 + O(x_1^2))\dot{x}_1 = g_1(x_1, y_2, c_1, c_2), \quad (56)$$

$$\dot{y}_2 = g_2(x_1, y_2, c_2), \quad (57)$$

where,

$$\begin{aligned} g_1(x_1, y_2, c_1, c_2) = & G_1(x_1^* + x_1 + w(y_2)O(x_1^2), y_1^* + O(x_1^2), c_1) \\ & + \nu H_1^s(x_1^* + x_1 + w(y_2)O(x_1^2), x_2^* + v(y_2)x_1 + O(x_1^2), y_1^* + O(x_1^2), y_2) \\ & - \frac{dy_1^*}{dy_2} \left( G_2(x_2^* + v(y_2)x_1 + O(x_1^2)) + \nu H_2^s(x_1^* + x_1 + w(y_2)O(x_1^2), \right. \\ & \left. x_2^* + v(y_2)x_1 + O(x_1^2), y_1^* + O(x_1^2), y_2) \right) \end{aligned} \quad (58)$$

$$\begin{aligned} g_2(x_1, y_2, c_2) = & G_2(x_2^* + v(y_2)x_1 + O(x_1^2)) + \nu H_2^s(x_1^* + x_1 + w(y_2)O(x_1^2), \\ & x_2^* + v(y_2)x_1 + O(x_1^2), y_1^* + O(x_1^2), y_2), \end{aligned} \quad (59)$$

and the desingularized system

$$\dot{x}_1 = g_1(x_1, y_2, c_1, c_2), \quad (60)$$

$$\dot{y}_2 = g_2(x_1, y_2, c_2)(2Kx_1 + O(x_1^2)). \quad (61)$$

### 3.3. Folded singularities of type FSN II and the existence of MMOs

As discussed in Section 2.2 a well known mechanism of transition to MMOs is Folded Saddle-Node of type II (FSN II), see [11], [19]. This is a codimension one transition, corresponding to the passage of the system from a parameter region with an excitable equilibrium to a parameter region of MMO dynamics. It can be described as follows in the context of system (56)-(57): as the regular parameter is varied a stable equilibrium of (56)-(57) approaches the fold, and, for the critical value of the regular parameter, satisfies the conditions:

$$g_1(0, y_2, c_1, c_2) = 0 \quad (62)$$

$$g_2(0, y_2, c_2) = 0. \quad (63)$$

On the other side of criticality there are MMOs as well as an equilibrium, now unstable.

In the following, we assume that the parameter  $c_2$  is fixed. Note that for  $\nu = 0$ ,  $y_1^*$  does not depend on  $y_2$  and thus  $dy_1^*/dy_2 = O(\nu)$ . We assume that, for  $\nu = 0$ , the equation  $G_2(x_2^*, y_2, c_2) = 0$  has a unique solution  $x_2^* = \psi_2(y_2, c_2)$  which is a stable equilibrium of the uncoupled  $(x_2, y_2)$  subsystem, and that  $G_{2,y_2} \neq 0$  at this point. By the implicit function theorem this gives a unique solution  $y_2$  for the equation (63), for  $\nu$  small enough. Recall that, by hypothesis, in the case  $\nu = 0$  we assume that, each uncoupled subsystem (39)-(41) and (40)-(42) admits a unique stationary point, one attractive on the stable manifold and one repulsive near the fold. This correspond to the nullcline configuration shown in Figure 1. Let us denote this particular value  $\bar{y}_2$ . It follows that this value  $\bar{y}_2$  determine particular values of




**Figure 1.** The  $\nu = 0$  limit configurations. Figure a) The uncoupled system (39)-(41) admits a unique repulsive stationary point near the fold. Figure b) The uncoupled system (40)-(42) admits a unique attractive stationary point on the stable manifold.

$(x_1^*, y_1^*, x_2^*) = (\bar{x}_1^*, \bar{y}_1^*, \bar{x}_2^*)$ . We assume that solving equation (62) with these values gives a unique value of  $c_1 = \bar{c}_1$ .

Recall from the discussion in Section 2.2 that a non-degenerate FSN II singularity corresponds to a transcritical bifurcation for system (60)-(61). We will show that such a transcritical bifurcation exists under some non-degeneracy assumptions on (35)-(38). Let us recall the general definition of a transcritical bifurcation, see [9, 5, 13] :

**Definition 1** *Let*

$$\dot{x} = X(x, c) \tag{64}$$

*be a system of differential equations in  $\mathbb{R}^n$  depending on a real parameter  $c$ . Let us assume that  $(\bar{x}, \bar{c})$  is a stationary point of (64):  $X(\bar{x}, \bar{c}) = 0$ . We say that this system admits a transcritical bifurcation at  $(\bar{x}, \bar{c})$  if there exists an open neighborhood  $U$  of  $x$  in  $\mathbb{R}^n$ , an open neighborhood  $V$  of  $c$  in  $\mathbb{R}$ , and  $\lambda_1, \lambda_2 : V \mapsto \mathbb{R}, \xi_1, \xi_2 : V \mapsto \mathbb{R}^n$  functions of class  $C^1$  such that:*

- (i)  $D_x X(\bar{x}, \bar{c})$  admits an eigenvalue equal to zero. This eigenvalue is simple and there is no other eigenvalue with real part equal to zero.
- (ii) For all  $c \in V$ ,  $\xi_1, \xi_2$  are the only stationary points of (64) in  $U$ , furthermore we have:  $\xi_i(\bar{c}) = \bar{x}$  and  $\xi_1 \neq \xi_2$  for  $c \neq \bar{c}$ .
- (iii) For all  $c \in V$ ,  $\lambda_i(c)$  is an eigenvalue of  $D_x X(\xi_i, c)$ , with  $\lambda_i(\bar{c}) = 0$  and  $\lambda_1 \lambda_2 < 0$  for  $c \neq \bar{c}$ .

The following proposition establishes the existence of a transcritical bifurcation for (60)-(61).

**Proposition 3** *Let us assume that, for system (60)-(61):*

$$\frac{dg_1}{dy_2} \frac{dG_1}{dx_1} \frac{dG_2}{dy_2}(0, \bar{y}_2, \bar{c}_1) < 0 (\text{resp } > 0), \tag{65}$$

then for  $c_1$  in a neighborhood of  $c_1 = \bar{c}_1$  and  $\nu$  small enough, there exists two stationary points for system (60)-(61):  $(x_{1,e}, y_{2,e})$  and  $(0, y_{2,fold})$ . Furthermore, if we assume that

$$\frac{dx_{1,e}}{dc_1} \frac{dg_2(0, y_{2,fold}(c_1))}{dc_1} > 0 (\text{resp } < 0), \quad (66)$$

then, for  $\nu$  sufficiently small, there is a transcritical bifurcation. It follows that  $(0, \bar{y}_2)$ , is an FSNII point.

### Proof

The folded stationary point,  $(0, y_{2,fold})$  is obtained by solving the equation  $g_1(0, y_2, c_1, c_2) = 0$ , i.e,  $g_1 = 0, x_1 = 0$ . For this, we apply the implicit function theorem to  $g_1(0, y_2, c_1, c_2)$  as a function of  $y_2$  and  $c_1$ . For  $y_2 = \bar{y}_2$  and  $c_1 = \bar{c}_1$ , we have  $g_1 = 0$ . From hypothesis (65), it follows that  $\frac{dg_1}{dy_2}(0, \bar{y}_2, \bar{c}_1, c_2) \neq 0$ , then the implicit function theorem gives the existence of a stationary point  $(x_1 = 0, y_2 = y_{2,fold})$  for  $c_1$  in a neighborhood of  $\bar{c}_1$ . The second stationary point,  $(x_{1,e}, y_{2,e})$ , is obtained by solving the equation  $g_1(x_1, y_2, c_1) = 0, g_2(x_1, y_2) = 0$ . Note that this second stationary point corresponds to a true stationary point of system (35)-(38). For  $\nu = 0$ , we know that these equations admit a unique solution. Using hypothesis (65), we can apply the implicit function in a neighborhood of  $\nu = 0$ . This gives the existence of a unique stationary point for  $\nu$  sufficiently small. For the first stationary point, the folded one  $(0, y_{2,fold})$  the jacobian is given by

$$J = \begin{pmatrix} G_{1,x_1} + O(\nu) & \frac{dg_1}{dy_2} \\ 2Kg_2 & 0 \end{pmatrix}. \quad (67)$$

This gives:

$$\det J = \left( \frac{dg_1}{dy_2} \right) 2Kg_2.$$

For the second stationary point, the ordinary one, the jacobian is given by:

$$\begin{pmatrix} G_{1,x_1} + O(\nu) + O(x_1) & \frac{dg_1}{dy_2} \\ KG_{2,x_1}x_1 + O(x_1^2) & KG_{2,y_2}x_1 + O(x_1^2) \end{pmatrix}. \quad (68)$$

This gives:

$$\det J = KG_{1,x_1}G_{2,y_2}x_1 + O(x_1\nu).$$

From hypothesis (66), it follows that system (60)-(61) admits a transcritical bifurcation at point  $(0, \bar{y}_2)$  when  $c_1$  crosses the value  $\bar{c}_1$ .

## 4. Example – coupled FitzHugh-Nagumo systems.

### 4.1. Simple fold line

We consider the following system

$$\begin{cases} \epsilon \dot{x}_1 &= F(x_1) - y_1 + \alpha_1(x_2 - x_1) \\ \epsilon \dot{x}_2 &= F(x_2) - y_2 + \alpha_2(x_1 - x_2) \\ \dot{y}_1 &= x_1 - c_1 + \beta_1(y_2 - y_1) \\ \dot{y}_2 &= x_2 - c_2 + \beta_2(y_1 - y_2) \end{cases} \quad (69)$$

with

$$F(z) = -z^3 + 3z.$$

We define

$$l(x_1, y_1, x_2, y_2) = \begin{cases} F(x_1) - y_1 + \alpha_1(x_2 - x_1) \\ F(x_2) - y_2 + \alpha_2(x_1 - x_2) \end{cases} \quad (70)$$

Points on the fold line are defined by the condition

$$\begin{cases} \det(Dl(x, y)) &= 0 \\ l_1(x, y) &= 0 \\ l_2(x, y) &= 0 \end{cases} \quad (71)$$

wich, for system (69), is given by

$$\begin{cases} (F'(x_1) - \alpha_1)(F'(x_2) - \alpha_2) - \alpha_1\alpha_2 &= 0 \\ y_1 &= F(x_1) + \alpha_1(x_2 - x_1) \\ y_2 &= F(x_2) + \alpha_2(x_1 - x_2). \end{cases}$$

Thanks to proposition 2, the existence of a *simple* fold line follows from  $F'(x_1) = 0$ ,  $F'(x_2) \neq 0$  and  $F''(x_1) \neq 0$  (the alternative choice would be  $F'(x_2) = 0$ ,  $F'(x_1) \neq 0$  and  $F''(x_2) \neq 0$ ). We obtain a solution of system (4.1) with  $(x_1, x_2, y_1)$  given as function of  $y_2$ . We denote this solution by  $(x_1^*, x_2^*, y_1^*) = (\varphi_1(y_2), \varphi_2(y_2), \varphi_3(y_2))$ . Note that we have obtained a curve in  $\mathbb{R}^4$  parametrized par  $y_2$ , see 2 for a general statement.

#### 4.2. Folded singularities and their nature

System (46)-(49) in context of (69) becomes

$$\begin{cases} \epsilon \dot{x}_1 &= -y_1 + f_1(x_1, y_2) + \alpha_1(x_2 - x_1) + O(\epsilon) \\ \epsilon \dot{x}_2 &= f_2(x_1, y_2) + \alpha_2(x_1 - x_2) + O(\epsilon) \\ \dot{y}_1 &= x_1^* + x_1 - c_1 + \beta_1(y_2 - (y_1^* + y_1)) \\ &\quad - \frac{dy_1^*}{dy_2}(x_2^* + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2)) \\ \dot{y}_2 &= x_2^* + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2) \end{cases} \quad (72)$$

where

$$f_1(x_1) = F'(x_1^*)x_1 + F''(x_1^*)\frac{x_1^2}{2} - x_1^3$$

and

$$f_2(x_2) = F'(x_2^*)x_2 + F''(x_2^*)\frac{x_2^2}{2} - x_2^3.$$

As in section 3.2, we diagonalize the linear part of the fast system. The eigenvalues of  $Df(x_1^*, x_2^*)$  are:

$$0 \text{ and } \lambda(y_2) = F'(x_1^*) + F'(x_2^*) - \alpha_1 - \alpha_2.$$

As explained in section (3.3), we focus on dynamics deccribed by configuration in Figure 1, thus  $F'(x_2^*) < 0$ . The associated eigenvectors are, respectively,

$$P_1 = \begin{pmatrix} 1 \\ \frac{\alpha_2}{F'(x_2^*) - \alpha_2} \end{pmatrix} \text{ and } P_2 = \begin{pmatrix} \frac{\alpha_1}{\alpha_2 - F'(x_2^*)} \\ 1 \end{pmatrix}.$$

We now diagonalize the fast subsystem using the transformation

$$X = P\tilde{X} \text{ with } X = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

where  $P$  is the matrix whose columns are the eigenvectors. We introduce the following notations:

$$w = \frac{\alpha_1}{\alpha_2 - F'(x_2^*)}, \quad v = \frac{\alpha_2}{F'(x_2^*) - \alpha_2}, \quad d = \frac{1}{1 - vw}.$$

After some transformations, system (72) becomes:

$$\begin{cases} \epsilon \dot{x}_1 &= d((F''(x_1^*)\frac{(x_1 + wx_2)^2}{2} - (x_1 + wx_2)^3 - y_1) \\ &\quad - w(F''(x_2^*)\frac{(vx_1 + x_2)^2}{2} - (vx_1 + x_2)^3)) + O(\epsilon) \\ \epsilon \dot{x}_2 &= \lambda(y_2)x_2 + d(-v(F''(x_1^*)\frac{(x_1 + wx_2)^2}{2} - (x_1 + wx_2)^3 - y_1) \\ &\quad + F''(x_2^*)\frac{(vx_1 + x_2)^2}{2} - (vx_1 + x_2)^3) + O(\epsilon) \\ \dot{y}_1 &= x_1^* + x_1 + wx_2 - c_1 + \beta_1(y_2 - (y_1^* + y_1)) \\ &\quad - \frac{dy_1^*}{dy_2}(x_2^* + bx_1 + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2)) \\ \dot{y}_2 &= x_2^* + vx_1 + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2). \end{cases}$$

Hence, the reduced equation is given by

$$0 = d((F''(x_1^*)\frac{(x_1 + wx_2)^2}{2} - (x_1 + wx_2)^3 - y_1) \quad (73)$$

$$- w(F''(x_2^*)\frac{(vx_1 + x_2)^2}{2} - (vx_1 + x_2)^3))$$

$$0 = \lambda(y_2)x_2 + d(-v(F''(x_1^*)\frac{(x_1 + wx_2)^2}{2} - (x_1 + wx_2)^3 - y_1) \quad (74)$$

$$+ F''(x_2^*)\frac{(vx_1 + x_2)^2}{2} - (vx_1 + x_2)^3)$$

$$\dot{y}_1 = x_1^* + x_1 + wx_2 - c_1 + \beta_1(y_2 - (y_1^* + y_1)) \quad (75)$$

$$- \frac{dy_1^*}{dy_2}(x_2^* + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2))$$

$$\dot{y}_2 = x_2^* + vx_1 + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2) \quad (76)$$

Now, using equation (74) and the implicit function theorem in (75), as  $\lambda(y_2) < 0$ , one can obtain  $x_2$  as a function of  $x_1, y_2$  which leads to,

$$x_2 = O(x_1^2).$$

Now, (74) can be rewritten in the form

$$y_1 = \frac{K}{2}x_1^2 + O(x_1^3) \quad (77)$$

with

$$\begin{aligned} K &= F''(x_1^*) - wv^2 F''(x_2^*) \\ &= F''(x_1^*) + O(\nu^3), \end{aligned}$$

where  $0 \leq \nu \leq \max(\alpha_1, \beta_1, \alpha_2, \beta_2)$ . Subsequently, we derivate (77), and plug into (75) and (76). We obtain,

$$\begin{aligned} (Kx_1 + O(x_1^2))\dot{x}_1 &= x_1^* + x_1 - c_1 + \beta_1(y_2 - y_1^*) \\ &\quad - \frac{dy_1^*}{dy_2}(x_2^* + bx_1 - c_2 + \beta_2(y_1^* - y_2)) + O(x_1^2) \\ \dot{y}_2 &= x_2^* + bx_1 - c_2 + \beta_2(y_1^* - y_2) + O(x_1^2) \end{aligned}$$

This gives the following desingularized system:

$$\dot{x}_1 = x_1^* + x_1 - c_1 + \beta_1(y_2 - y_1^*) \quad (78)$$

$$- \frac{dy_1^*}{dy_2}(x_2^* + bx_1 - c_2 + \beta_2(y_1^* - y_2)) + O(x_1^2) \quad (79)$$

$$\dot{y}_2 = (x_2^* + bx_1 - c_2 + \beta_2(y_1^* - y_2) + O(x_1^2))(Kx_1 + O(x_1^2)) \quad (80)$$

$$(81)$$

Hence using notations of section 3.3, we have:

$$g_1(x_1, y_2) = x_1^* + x_1 - c_1 + \beta_1(y_2 - y_1^*) - \frac{\partial y_1^*}{\partial y_2}(x_2^* + bx_1 - c_2 + \beta_2(y_1^* - y_2)) + O(x_1^2)$$

and

$$g_2(x_1, y_2) = x_2^* + bx_1 - c_2 + \beta_2(y_1^* - y_2) + O(x_1^2).$$

**Proposition 4** *Let us assume  $c_2 \ll -1$  and  $\alpha_1 \leq \beta_1$ . For  $\alpha_1, \alpha_2, \beta_1, \beta_2$  small enough, let  $\bar{y}_2$  be the solution of  $g_2(0, y_2) = 0$  and define  $\bar{c}_1 = x_1^* + \beta_1(\bar{y}_2 - y_1^*)$ . Then, for  $\alpha_1, \alpha_2, \beta_1, \beta_2$  in a small neighborhood of zero, the system (80)-(81), admits a transcritical bifurcation at point  $(x_1 = 0, y_2 = \bar{y}_2, c_1 = \bar{c}_1)$ . As the parameter  $c_1$  crosses the value  $\bar{c}_1$  from left to right, the folded stationary point passes from a folded saddle to a repulsive folded node, whereas the ordinary stationary point passes from a repulsive node to a saddle.*

### Proof

This proposition is a specific case of proposition (3). We will verify that:

$$\frac{dg_1}{dy_2}(0, \bar{y}_2, \bar{c}_1) > 0, \quad (82)$$

$$\frac{dG_1}{dx_1} \frac{dG_2}{dy_2} < 0, \quad (83)$$

$$\frac{dx_{1,e}}{dc_1} > 0, \quad (84)$$

$$\frac{dg_2(0, \bar{y}_{2,fold}(c_1))}{dc_1} > 0. \quad (85)$$

We first show the existence of two distinct stationary points for system (80)-(81). We start with hypothesis (82). Here, this hypothesis reads as,

$$\beta_1 - \frac{\alpha_1}{9(1 - \bar{x}_2^{*2})^2} + O(\nu^2) > 0$$

It is sufficient to have:

$$\bar{x}_2^{*2} \notin ]1 - \sqrt{\frac{\alpha_1}{9\beta_1}}, 1 + \sqrt{\frac{\alpha_1}{9\beta_1}}[$$

This holds since  $\bar{x}_2^* = c_2 + O(\nu)$ ,  $c_2 \ll -1$  and  $\alpha_1 \leq \beta_1$ . Now, we deal with hypothesis (83). Here, it reads:

$$\frac{dx_2^*}{dy_2} < 0. \quad (86)$$

This holds since  $\frac{dx_2^*}{dy_2} = \frac{1}{F'(x_2)} + O(\nu) < 0$  for  $\nu$  sufficiently small. Hypothesis (84) is verified since we have  $\frac{dx_{1,e}}{dc_1} + O(\nu) > 0$  or  $\nu$  sufficiently small.

Finally, we come to hypothesis (85). From (82), we know that  $\frac{dg_1}{dy_2}(0, \bar{y}_2, \bar{c}_1) > 0$ . From (83), we know that  $\frac{dg_2}{dy_2} < 0$ . Since we have,  $\frac{dg_1}{dc_1} < 0$ , it follows that  $\frac{dy_2}{dc_1} < 0$ , and then  $\frac{dg_2}{dc_1} < 0$ .

## 5. Numerical simulations

We proceed numerical integration of system (69) on time interval  $[0, 500]$ , using a Runge-Kutta 4 method with time step of 0.001. The parameter values are the following  $\varepsilon = 0.1, \alpha_1 = \beta_1 = \alpha_2 = \beta_2 = 0.01, c_2 = -1.5$ . The parameter  $c_1$  is varied in a small neighborhood around the value  $-0.99$ . In figure 2, we represent the results of the simulation when  $c_1 = -0.99$ . Figure 2,a) represents the time-evolution of the variable  $x_1$ . Figure 2,b) represents the behaviour of the system in the  $(x_1, y_1)$  phase-plane. It shows, that the system evolves asymptotically toward a stable stationary point. In figure 3, we represent the results of the simulation when  $c_1 = -0.9883$ . Figure 3,a) represents the time-evolution of the variable  $x_1$ . Figure 3,b) represents the behaviour of the system in the  $(x_1, y_1)$  phase-plane. It shows that the system evolves asymptotically toward a limit cycle of small amplitude. In figure 3, we represent the results of the simulation when  $c_1 = -0.988295$ . Figure 4,a) represents the time-evolution of the variable  $x_1$ . Figure 4,b) represents the time-evolution of the variable  $x_2$ . Figure 4,c) represents the behaviour of the system in the  $(x_1, y_1)$  phase space. Figure 4,d) represents the behaviour of the system in the  $(x_2, y_2)$  phase space. It shows the emergence of the MMO's. Figure 5,d) represents the evolution of the system in the  $(x_2, y_2)$  phase space. Figure 5 represents the reduced manifold:  $(x_1, x_2, y_1 = F(x_1) + \alpha_1(x_2 - x_1))$ , and the behaviour of the system in the  $(x_1, x_2, y_1)$  phase-space. It shows the emergence of the MMO's.




**Figure 2.** Simulation of system (69) for parameter  $c_1 = -0.99$ . Figure a) The time-evolution of  $x_1$ . Figure b) The behaviour of the system in the  $(x_1, y_1)$  phase-plane. It shows that the system evolves asymptotically toward its stable stationary point.




**Figure 3.** Simulation of system (69) for parameter  $c_1 = -0.9883$ . Figure a) The time-evolution of  $x_1$ . Figure b) The behaviour of the system in the  $(x_1, y_1)$  phase-plane. It shows that the system evolves asymptotically toward a limit cycle of small amplitude.

## Acknowledgments

We would like to thank: Région Haute Normandie and - FEDER (RISC project) for financial support.




**Figure 4.** Simulation of system (69) for parameter  $c_1 = -988295$ . Figure a) The time-evolution of  $x_1$ . Figure b) The time-evolution of  $x_2$ . Figure c) The behaviour of the system in the  $(x_1, y_1)$  phase-plane. The behaviour of the system in the  $(x_2, y_2)$  phase space. It shows the emergence of MMO's.




**Figure 5.** Simulation of system (69) for parameter  $c_1 = -988295$ . The reduced manifold:  $(x_1, x_2, y_1 = F(x_1) + \alpha_1(x_2 - x_1))$  and the behaviour of the system in the  $(x_1, x_2, y_1)$  phase-space (green curve). It shows the emergence of MMO's.

- [1] E. Benoît. Canards et enlacements. *Publ. Math. IHES*, 72:63–91, 1990.
- [2] E. Benoît, J.-L. Callot, F. Diener, and M. Diener. Chasse au canard. *Collect. Math.*, 31:37–119, 1981.
- [3] M. Brøns, M. Krupa, and M. Wechselberger. Mixed-Mode Oscillations Due to Generalized Canard Phenomenon. In *Bifurcation Theory and spatio-temporal pattern formation*, pages 39– 64. Fields Institute Communications, 2006.
- [4] M. Brøns. Bifurcations and instabilities in the Greitzer model for compressor system surge. *Math. Eng. Ind.*, 2(1):51–63, 1988.
- [5] J.P. Francoise. *Oscillations en biologie*. Springer, 2005.
- [6] R. Curtu and J. Rubin. Interaction of Canard and Singular Hopf Mechanisms in a Neural Model. *SIAM J. Appl. Dyn. Syst.*, 10 (4):1443–1479, 2011.
- [7] M. Desroches, J. Guckenheimer, B. Krauskopf, C. Kuehn, H. M Osinga, and M. Wechselberger. Mixed-mode oscillations with multiple time scales. *SIAM Review*, 54(2):211–288, 2012.
- [8] B. Ermentrout and M. Wechselberger. Canards, Clusters and Synchronization in a Weakly Coupled Interneuron Model. *SIAM J. Appl. Dyn. Syst.*, 8(1):253–278, 2009.
- [9] J. Guckenheimer and P. Holmes. *Non linear oscillations, dynamical systems and bifurcations of vector fields*. Springer-Verlag, 1985.
- [10] J. Guckenheimer. Singular Hopf bifurcation in systems with two slow variables. *SIAM J. Appl. Dyn. Syst.*, 7:1355–1377, 2008.
- [11] M. Krupa and M. Wechselberger. Local analysis near a folded saddle-node singularity. *J. Differential Equations*, 248:2841–2888, 2010.
- [12] A. Milik, P. Szmolyan, H. Loeffelmann, and E. Groeller. Geometry of mixed-mode oscillations in the 3d autocatalator. *Int. J. Bifur. Chaos*, 8:505–519, 1998.
- [13] A. Otto. *Étude de quelques modèles épidémiologiques par les méthodes de calcul formel , in french*. PhD thesis, Université de Rennes, 2009.
- [14] H. Rotstein, M. Wechselberger, and N. Kopell. Canard Induced Mixed-Mode Oscillations in a Medial Entorhinal Cortex Layer II Stellate Cell Model. *SIAM J. Appl. Dyn. Syst.*, 7(4):1582–1611, 2008.
- [15] J. Rubin and M. Wechselberger. Giant squid - hidden canard: the 3d geometry of the Hodgkin-Huxley model. *Biol. Cyber.*, 97(5), 2007.
- [16] P. Szmolyan and M. Wechselberger. Canards in  $\mathbb{R}^3$ . *J. Differential Equations*, 177:419–453, 2001.
- [17] T. Vo, R. Bertram, J. Tabak, and M. Wechselberger. Mixed mode oscillations as a mechanism for pseudo-plateau bursting. *J. Comput. Neurosci.*, 28 (3):443–458, 2010.
- [18] M. Wechselberger. Existence and Bifurcations of Canards in  $\mathbb{R}^3$  in the Case of the Folded Node. *SIAM J. Appl. Dyn. Syst.*, 4:101–139, 2005.
- [19] M. Wechselberger. Canards. 2(4):1356, 2007.
- [20] M. Wechselberger. A propos de canards (apropos canards). *Trans. Amer. Math. Soc.*, 364(6):3289–3309, 2011.